

## Math 131 PracTest 2

0. Finish Lab 8, problems 1–10 except 6. Review Labs 6 and 7—answers are online.

1. Let  $S$  be the region enclosed by the  $y$  axis,  $y = x^2 + 4$ , and  $y = 2x^2$  in the first quadrant only. Sketch the region.

a) Rotate  $S$  about the  $x$ -axis and find the resulting volume. (Ans:  $512\pi/15$ )



b) Rotate  $S$  about the  $y$ -axis and find the resulting volume. (Ans:  $8\pi$ )

2. Sketch the region  $R$  in the first quadrant enclosed by the  $x$ -axis,  $y = \sqrt{x}$  and  $y = x - 2$ .

a) Rotate  $R$  about the  $x$ -axis and find the volume. (Ans:  $16\pi/3$ )

b) Rotate  $R$  about the  $y$ -axis and find the volume. Try both methods. (Ans:  $184\pi/15$ )

3. A wooden doorstop with right triangular cross-sections is 20 cm long and 5 cm high at its tall end and 4 cm wide. Find its volume. Hint: Find the equation of the line that forms the top edges and use similar triangles to find the cross-sectional area. (Ans:  $\frac{200}{3}$  cm<sup>3</sup>.)



4. a) Let  $R$  be the region between  $y = \ln x$  and the  $x$  axis on the interval  $[1, e^2]$ . Rotate  $R$  around the  $x$  axis and find the resulting volume. What integration technique should you use? (Answer:  $\pi(2e^2 - 1)$ .)

b) Rotate  $R$  about the  $y$  axis and find the volume using the shell method. (Answer:  $\frac{\pi}{2}(1 + 3e^4)$ .)

c) Rotate  $R$  about the  $y$  axis and find the volume using the disk method. (Answer:  $\frac{\pi}{2}(1 + 3e^4)$ .)

5. Let  $R$  be the region between  $y = \sin \pi x$  and the  $x$  axis on the interval  $[0, 1]$ . Rotate  $R$  about the  $x$ -axis and find the resulting volume. Hint: Use an identity. (Answer:  $\pi/2$ .)

6. Find the average value of  $f(x) = \sin^3 x$  on the interval  $[0, \pi]$ . [Ans:  $4/3\pi$ .]

7. Find the volume of the solid region generated when the area in the first quadrant enclosed by  $y = \cos x$ ,  $y = 0$ ,  $x = 0$  and  $x = \pi/2$  is revolved around the  $y$ -axis. Use shells. [Ans:  $\pi^2 - 2\pi$ .]

8. a) Find  $\int_0^{\pi/4} x \tan^2 x \, dx$ . [Ans:  $\frac{\pi}{4} - \ln \sqrt{2} - \frac{\pi^2}{32}$ .]

b) Let  $R$  be the region between  $y = \tan^2 x$ ,  $x = \pi/4$ , and the  $x$ -axis in the first quadrant. Rotate  $R$  about the  $y$  axis and find the volume using the shell method. Re-use part (a). [Ans:  $\frac{\pi^2}{2} - 2\pi \ln \sqrt{2} - \frac{\pi^3}{16}$ .]

c) Let  $R$  be the region enclosed by  $y = \tan^{-1} x$ , the  $y$ -axis, and  $y = \pi/4$  in the first quadrant. Rotate  $R$  about the  $y$ -axis to form a tank. If it is full of a liquid whose density is 64 lbs/cu. ft., how much work is lost if it leaks out the bottom and drops to ground level? Hint: Use part (a)! [Ans:  $-16\pi^2 + 64\pi \ln \sqrt{2} + 2\pi^3$ .]

9. Find the length of  $f(x) = \frac{4}{3}x^{3/2} + 1$  on the interval  $[0, 2]$ . (Answer:  $13/3$ )

10. Let  $f(x) = \frac{1}{6}x^3 + \frac{1}{2}x^{-1}$  on  $[2, 3]$ . Find the arc length. (Answer:  $13/4$ .)




11. Find the arc length of  $f(x) = \ln \cos x$  on  $[0, \pi/4]$ . (Answer:  $\ln |\sqrt{2} + 1|$ .)

12. Find the volume of the solid region generated when the area in the first quadrant enclosed by  $y = \sqrt{\frac{x}{9+x^2}}$ ,  $y = 0$ , and  $x = 2$  is revolved around the  $x$ -axis. [Ans:  $\frac{\pi}{2}[\ln \frac{13}{9}]$ ]

13. Find the volume of the solid region generated when the area in the first quadrant enclosed by  $y = \cos x$ ,  $y = 0$ ,  $x = 0$  and  $x = \pi/2$  is revolved around the  $y$ -axis. Use shells. [Ans:  $\pi^2 - 2\pi$ .]

14. a) A small farm elevated water tank is in the shape obtained from rotating the region in the first quadrant enclosed by the curves  $y = 10 - \frac{1}{2}x^2$ ,  $y = 8$ , and the  $y$ -axis about the  $y$ -axis. Find the work “lost” if the water (62.5 lbs/ft<sup>3</sup>) leaks onto the ground from a hole in the bottom of the tank. (Answer:  $-6500\pi/3$  ft-lbs.)
- b) Find the work “lost” if the water leaks onto the ground from a hole in the side of the tank at height 9 feet. (Answer:  $-1750\pi/3$  ft-lbs.)
- c) Set up integral for the work to empty a tank containing just one foot of water over the top edge.

Quiz



1. **Rotation about the  $x$ -axis.** Let  $R$  be the entire region enclosed by  $y = x^2$  and  $y = 2 - x^2$  in the upper half-plane. Rotate  $R$  about the  $x$ -axis. The resulting volume is given by:

- a)  $\pi \int_{-1}^1 (x^2)^2 dx - \pi \int_{-1}^1 (2 - x^2)^2 dx$       b)  $\pi \int_0^1 (2 - x^2)^2 dx - \pi \int_0^1 (x^2)^2 dx$
- c)  $\pi \int_{-1}^1 (2 - x^2)^2 dx + \pi \int_{-1}^1 (x^2)^2 dx$       d)  $\pi \int_{-1}^1 (2 - x^2)^2 dx - \pi \int_{-1}^1 (x^2)^2 dx$
- e)  $\pi \int_0^1 (\sqrt{y})^2 dy + \pi \int_1^2 (\sqrt{2 - y})^2 dy$       f) None of these

2. **Rotation about the  $x$ -axis.** Let  $S$  be the region enclosed by the  $x$ -axis,  $y = \sqrt{x}$ , and  $y = 2 - x$ . The volume generated by revolving  $S$  about the  $x$ -axis is:

- a)  $\pi \int_0^1 (\sqrt{x})^2 dx - \pi \int_1^2 (2 - x)^2 dx$       b)  $2\pi \int_0^1 x(2 - x) dx - 2\pi \int_0^1 x\sqrt{x} dx$
- c)  $\pi \int_0^1 (\sqrt{x})^2 dx + \pi \int_1^2 (2 - x)^2 dx$       d)  $\pi \int_0^1 (2 - y)^2 - (y^2)^2 dy$
- e)  $\pi \int_0^2 (\sqrt{x})^2 dx - \pi \int_1^2 (2 - x)^2 dx$       f) None of these

3. **Rotation about the  $y$ -axis.** Let  $T$  be the region enclosed by the  $y$ -axis,  $y = \sqrt{x}$ , and  $y = 2 - x$  (a different region than in Problem 2). The volume generated by revolving  $T$  about the  $y$ -axis is:

- a)  $\pi \int_0^2 (y^2)^2 dy - \pi \int_0^2 (2 - y)^2 dy$       b)  $2\pi \int_0^1 x(2 - x)^2 dx - 2\pi \int_0^1 x(\sqrt{x})^2 dx$
- c)  $\pi \int_0^1 (\sqrt{x})^2 dx + \pi \int_1^2 (2 - x)^2 dx$       d)  $\pi \int_0^1 (2 - y)^2 dy + \pi \int_1^2 (y^2)^2 dy$
- e)  $2\pi \int_0^1 x(2 - x) dx - 2\pi \int_0^1 x(\sqrt{x}) dx$       f) None of these

4. **Rotation about the  $y$ -axis.** Let  $U$  be the region enclosed by the  $y$ -axis, the  $x$ -axis,  $y = x - 2$ , and  $y = 4 - x$ . The volume generated by revolving  $U$  about the  $y$ -axis is:

- a)  $\pi \int_0^1 (y - 2)^2 + \pi \int_1^4 (4 - y)^2 dy$       b)  $2\pi \int_0^3 x(4 - x)^2 dx - 2\pi \int_2^3 x(x - 2)^2 dx$
- c)  $2\pi \int_0^4 x(4 - x) dx - 2\pi \int_2^3 x(x - 2) dx$       d) a and c
- e) a and b      f) None of these

5. Integral Mix Up: Gotta game plan?

$$\begin{array}{lll} \text{a)} \int (4x^3 + 1) \ln x \, dx & \text{b)} \int x e^{x+1} \, dx & \text{c)} \int e^{2x} \cos x \, dx \\ \text{d)} \int x \sec^2 x \, dx & \text{e)} \int \cos^3(4x) \, dx & \\ \text{f)} \int \ln(2x^3) \, dx & \text{g)} \int \sin^2(5x) \, dx & \text{h)} \int x^2 \ln x^2 \, dx \\ \text{i)} \int \arctan 2x \, dx & \text{j)} \int 2 \sec^3 x \tan x \, dx \text{ See hint} & \text{k)} \int x \sin^2 x \, dx \quad \text{l)} \int \tan^3 x \, dx \end{array}$$

Hint:  $2 \sec^3 x \tan x = 2 \sec^2 x \sec x \tan x$

6. Determine  $\int 2x \arcsin x \, dx$ . You will need to use several different methods.

7. Draw the right triangle associated with each of these square roots and label the sides. For each, solve for  $u$ ,  $du$ , and the given square root in terms of an angle  $\theta$ .

$$\text{a)} \sqrt{u^2 - a^2} \quad \text{b)} \sqrt{a^2 - u^2} \quad \text{c)} \sqrt{u^2 + a^2}$$

8. Determine these integrals. *Caution:* A variety of techniques are required. Where necessary, make use of the triangles that you just drew.

$$\begin{array}{lll} \text{a)} \int \frac{\sqrt{x^2 - 1}}{x} \, dx & \text{b)} \int \frac{1}{(36 + x^2)^{3/2}} \, dx & \text{c)} \int \frac{1}{x^2 \sqrt{x^2 - 4}} \, dx \\ \text{d)} \int x \sqrt{4 - x^2} \, dx & \text{e)} \int \frac{4}{4 - x^2} \, dx & \text{f)} \int \frac{x^3}{\sqrt{4 - x^2}} \, dx \quad \text{g)} \int \frac{1}{\sqrt{9 - 4x^2}} \, dx \end{array}$$

9. Determine  $\int \frac{1}{9x^2 \sqrt{1 - 9x^2}} \, dx$ .

10. More Integral Mix Up: Before working these out, go through and classify each by the technique that you think will apply: substitution, parts, parts twice, trig methods, or ordinary methods.


$$\begin{array}{lll} \text{a)} \int x e^{2x} \, dx & \text{b)} \int x e^{x^2+1} \, dx & \text{c)} \int e^x \cos x \, dx \\ \text{d)} \int x^2 \sin x \, dx & \text{e)} \int \sin^4(2x) \, dx & \text{f)} \int x \cos(x^2) \, dx \quad \text{g)} \int \frac{(\ln x)^2}{x} \, dx \\ \text{h)} \int \ln(x^2) \, dx & \text{i)} \int \ln \sqrt{x} \, dx & \text{j)} \int x^2 \ln x \, dx \quad \text{k)} \int \cos^2(12x) \, dx \\ \text{l)} \int \tan x \, dx & \text{m)} \int \arctan x \, dx & \text{n)} \int \sin(5x) \cos(2x) \, dx \\ \text{o)} \int x \sqrt{4 - x^2} \, dx & \text{p)} \int \frac{1}{\sqrt{9 - 4x^2}} \, dx & \end{array}$$

## Math 131 Practest 2: Selected Answers

1. a)  $V = \pi \int_0^2 (x^2 + 4)^2 - (2x^2)^2 dx = \pi \int_0^2 (x^4 + 8x^2 + 16) - 4x^4 dx = \pi \int_0^2 8x^2 + 16 - 3x^4 dx = \pi \left[ \frac{8}{3}x^3 + 16x - \frac{3}{5}x^5 \right]_0^2 = \pi \left[ \left( \frac{64}{3} + 32 - \frac{96}{5} \right) - 0 \right] = 512\pi/15$

b)  $V = 2\pi \int_0^2 x(x^2 + 4) - x(2x^2) dx = 2\pi \int_0^2 (x^3 + 4x) - 2x^3 dx = 2\pi \int_0^2 4x - x^3 dx = 2\pi \left[ 2x^2 - \frac{1}{4}x^4 \right]_0^2 = 2\pi[(8 - 4)] = 8\pi$

2. Intersect:  $(\sqrt{x})^2 = (x - 2)^2 \Rightarrow x^2 - 5x + 4 = 0 \Rightarrow x = 4$  (not  $x = 1$ ). Note  $x = y^2$  and  $x = y + 2$



a)  $V = \pi \int_0^4 (x^{1/2})^2 dx - \pi \int_2^4 (x - 2)^2 dx = \pi \left[ \frac{1}{2}x^2 \right]_0^4 - \pi \left[ \frac{1}{3}(x - 2)^3 \right]_2^4 = \pi[(8 - 0) - (\frac{8}{3} - 0)] = 16\pi/3$

b)  $V = \pi \int_0^2 (y + 2)^2 - (y^2)^2 dy = \pi \left[ \frac{1}{3}(y + 2)^3 - \frac{1}{5}y^5 \right]_0^2 = \pi \left[ \left( \frac{64}{3} - \frac{32}{5} \right) - \left( \frac{8}{3} - 0 \right) \right] = 184\pi/15$

3. The line for the height is  $y = \frac{5}{20}x = \frac{1}{4}x$ . The line for the base width is  $y = \frac{4}{20}x = \frac{1}{5}x$ . So the cross-sectional area is  $A(x) = \frac{1}{2}bh = \frac{1}{2}(\frac{1}{5}x)(\frac{1}{4}x) = \frac{1}{40}x^2$ . So

$$V = \int_0^{20} \frac{1}{40}x^2 dx = \frac{1}{120}x^3 \Big|_0^{20} = \frac{200}{3} \text{ cm}^3.$$

4. a) Parts twice:

$u = (\ln x)^2$ $du = \frac{2 \ln x}{x} dx$	$dv = dx$ $v = x$	$\int (\ln x)^2 dx = x(\ln x)^2 - \int 2 \ln x dx$
$u = 2 \ln x$ $du = \frac{2}{x} dx$	$dv = dx$ $v = x$	$\int (\ln x)^2 x dx = x(\ln x)^2 - [2x \ln x - \int 2 dx] = x(\ln x)^2 - 2x \ln x + 2x + c$

So  $\pi \int_1^{e^2} (\ln x)^2 dx = \pi [x(\ln x)^2 - 2x \ln x + 2x]_1^{e^2} = \pi[(4e^2 - 4e^2 + 2e^2) - (0 - 0 + 1)] = \pi(2e^2 - 1)$ .

b) Parts:

$u = \ln x$ $du = \frac{1}{x} dx$	$dv = 2\pi x dx$ $v = \pi x^2$	$\int_1^{e^2} 2\pi x \ln x dx = \pi x^2 \ln x \Big _1^{e^2} - \int_1^{e^2} \pi x dx$ $= \pi x^2 \ln x - \frac{\pi}{2} x^2 \Big _1^{e^2} = (2\pi e^4 - \frac{\pi}{2} e^4) - (0 - \frac{\pi}{2}) = \frac{3\pi}{2} e^4 + \frac{\pi}{2}$
--------------------------------------	-----------------------------------	---

c)  $V = \pi \int_0^2 (e^2)^2 - (e^y)^2 dy = \pi \int_0^2 e^4 - e^{2y} dy = \pi [e^4 y - \frac{1}{2}e^{2y}]_0^2 = \pi[(2e^4 - \frac{1}{2}e^4) - (0 - \frac{1}{2})] = \frac{3\pi}{2} e^4 + \frac{\pi}{2}$ .

5.  $V = \pi \int_0^1 \sin^2(\pi x) dx = \pi \int_0^1 \frac{1}{2} - \frac{1}{2} \cos(2\pi x) dx = \pi \left[ \frac{1}{2}x - \frac{1}{4\pi} \sin(2\pi x) \right]_0^1 = \pi \left[ \left( \frac{1}{2} - 0 \right) - 0 \right] = \frac{\pi}{2}$

6. Via Reduction: Ave Val =  $\frac{1}{\pi - 0} \int_0^\pi \sin^3 x dx = \frac{1}{\pi} \left[ -\frac{1}{3} \sin x \cos x + \frac{2}{3} \int_0^\pi \sin x dx \right]$   
 $= \frac{1}{\pi} \left[ -\frac{1}{3} \sin x \cos x - \frac{2}{3} \cos x \right] \Big|_0^\pi \frac{1}{\pi} \left[ 0 - \left( -\frac{2}{3} \right) \right] - \left[ 0 - \frac{2}{3} \right] = \frac{4}{3\pi}$ .

7.  $V = \int_0^{\pi/2} 2\pi x \cos x dx = 2\pi [x \sin x - \int \sin x dx]_0^{\pi/2} = 2\pi [x \sin x + \cos x dx]_0^{\pi/2} = \pi^2 - 2\pi$ .

8. a) Trig id and then parts:  $\int_0^{\pi/4} x \tan^2 x dx = \int_0^{\pi/4} x(\sec^2 x - 1) dx = \int_0^{\pi/4} x \sec^2 x - x dx$ . To do  $\int x \sec^2 x dx$ :

$u = x$ $du = dx$	$dv = \sec^2 x dx$ $v = \tan x$	$\int x \sec^2 x dx = x \tan x - \int \tan x dx$ $\int x \sec^2 x dx = x \tan x - \ln  \sec x $
----------------------	------------------------------------	--

So  $\int_0^{\pi/4} x \tan^2 x dx = \int_0^{\pi/4} x \sec^2 x - x dx = [x \tan x - \ln |\sec x| - \frac{1}{2}x^2]_0^{\pi/4} = \frac{\pi}{4} - \ln \sqrt{2} - \frac{\pi^2}{32}$ .

b)  $V = \int_0^{\pi/4} 2\pi x \tan^2 x dx = 2\pi \left( \frac{\pi}{4} - \ln \sqrt{2} - \frac{\pi^2}{32} \right) = \frac{\pi^2}{2} - 2\pi \ln \sqrt{2} - \frac{\pi^3}{16} = \frac{\pi^2}{2} - \pi \ln 2 - \frac{\pi^3}{16}$ .

c) Note:  $x = \tan y$ , so  $W = 64 \int_0^{\pi/4} (0 - y)\pi \tan^2 y dy = -64\pi \left( \frac{\pi}{4} - \ln \sqrt{2} - \frac{\pi^2}{32} \right) = -16\pi^2 + 64\pi \ln \sqrt{2} + 2\pi^3$ .

9.  $AL = \int_0^2 \sqrt{1 + 4x} dx = \frac{1}{4} \cdot \frac{2}{3} (1 + 4x)^{3/2} \Big|_0^2 = \frac{1}{6} (27 - 1) = \frac{13}{3}$ .

$$10. AL = \int_2^3 \sqrt{1 + (\frac{1}{2}x^2 - \frac{1}{2}x^{-2})^2} dx = \int_2^3 \sqrt{(\frac{1}{2}x^2 + \frac{1}{2}x^{-2})^2} dx = \int_2^3 \frac{1}{2}x^2 + \frac{1}{2}x^{-2} dx = \frac{1}{6}x^3 - \frac{1}{2}x^{-1} \Big|_2^3 = \frac{13}{4}.$$

$$11. AL = \int_0^{\pi/4} \sqrt{1 + \tan^2 x} dx = \int_0^{\pi/4} \sqrt{\sec^2 x} dx = \int_0^{\pi/4} \sec x dx = \ln |\sec x + \tan x| \Big|_0^{\pi/4} = \ln |\sqrt{2} + 1|.$$

$$12. V = \pi \int_0^2 \left( \sqrt{\frac{x}{9+x^2}} \right)^2 dx = \pi \int_0^2 \frac{x}{9+x^2} dx = \frac{\pi}{2} \ln(9+x^2) \Big|_0^2 = \frac{\pi}{2} [\ln 13 - \ln 9].$$

$$13. V = \int_0^{\pi/2} 2\pi x \cos x dx = 2\pi [x \sin x - \int \sin x dx] \Big|_0^{\pi/2} = 2\pi [x \sin x + \cos x dx] \Big|_0^{\pi/2} = \pi^2 - 2\pi.$$

14. a) Save work! Since the radius of a cross-section of the tank is  $x$ , we need to solve for  $x^2$  (not  $x$ ): but  $y = 10 - \frac{1}{2}x^2$ , so  $\frac{1}{2}x^2 = 10 - y$  or  $x^2 = 20 - 2y$ .

$$\begin{aligned} W &= 62.5 \int_8^{10} \pi(20 - 2y)(0 - y) dy = 62.5\pi \int_8^{10} 2y^2 - 20y dy = 62.5\pi [2y^3/3 - 10y^2] \Big|_8^{10} \\ &= 62.5\pi [(2000/3 - 1000) - (1024/3 - 640)] = -6500\pi/3 \text{ lbf} \end{aligned}$$

b) Only the lower limit changes to 9 since the upper part of the barrel leaks out:

$$\begin{aligned} W &= 62.5 \int_9^{10} \pi(20 - 2y)(0 - y) dy = -62.5\pi [2y^3/3 - 10y^2] \Big|_9^{10} \\ &= -62.5\pi [(2000/3 - 1000) - (486 - 810)] = -1750\pi/3 \text{ lbf} \end{aligned}$$

c) Note the limits and the height 'moved to':  $W = 62.5 \int_8^9 \pi(20 - 2y)(10 - y) dy$

**Quiz Answers:** 1 (d) 2 (c) 3 (e) 4 (f)

5. a)	$u = \ln x \quad dv = 4x^3 + 1 dx$ $du = \frac{1}{x} dx \quad v = x^4 + x$	$\int (4x^3 + 1) \ln x dx = (x^4 + x) \ln x - \int \frac{x^4+x}{x} dx$ $= (x^4 + x) \ln x - \int x^3 + 1 dx = (x^4 + x) \ln x - \frac{1}{4}x^4 - x + c dx$
-------	---	---

b)	$u = x \quad dv = e^{x+1} dx$ $du = dx \quad v = e^{x+1}$	$\int x e^{x+1} dx = x e^{x+1} - \int e^{x+1} dx = x e^{x+1} - e^{x+1} + c = e^{x+1}(x - 1) + c$
----	--	--

c) Parts twice, "circle around". Note signs and constants:

$u = e^{2x} \quad dv = \cos x dx$ $du = 2e^{2x} dx \quad v = \sin x$	$\int e^{2x} \cos x dx = e^{2x} \sin x - \int 2e^{2x} \sin x dx$
$u = 2e^{2x} \quad dv = \sin x dx$ $du = 4e^{2x} dx \quad v = -\cos x$	$\int e^{2x} \sin x dx = e^{2x} \sin x - [-2e^{2x} \cos x + 4 \int e^{2x} \cos x dx]$ So, $5 \int e^{2x} \cos x dx = e^{2x} \sin x + 2e^{2x} \cos x + c$ Thus, $\int e^{2x} \cos x dx = \frac{1}{5} e^{2x} (\sin x + 2 \cos x) + c$

d)	$u = x \quad dv = \sec^2 x dx$ $du = dx \quad v = \tan x$	$\int x \sec^2 x dx = x \tan x - \int \tan x dx = x \tan x + \ln  \cos x  + c$
----	--	--

e) Reduction:  $u = 4x$  so  $\frac{1}{4} du = dx$ .  $\int \cos^3(4x) dx = \frac{1}{4} \int \cos^3 u du$

$$= \frac{1}{4} \left[ \frac{\cos^2 u \sin u}{3} + \frac{2}{3} \int \cos u du \right] = \frac{1}{4} \left[ \frac{\cos^2 u \sin u}{3} + \frac{2}{3} \sin u \right] = \frac{1}{12} \cos^2(4x) \sin(4x) + \frac{1}{6} \sin(4x) + c$$

f)	$u = \ln(2x^3) \quad dv = dx$ $du = \frac{6x^2}{2x^3} dx = \frac{3}{x} dx \quad v = x$	$\int \ln(2x^3) dx = x \ln(2x^3) - \int \frac{3x}{x} dx$ $= x \ln(2x^3) - \int 3 dx = x \ln(2x^3) - 3x + c$
----	---	--

g)  $\int \sin^2(5x) dx = \int \frac{1}{2} - \frac{1}{2} \cos(10x) dx = \frac{1}{2}x - \frac{1}{20} \sin(10x) + c.$

h)	$u = \ln(x^2)$	$dv = x^2 dx$	$\int x^2 \ln x^2 dx = \frac{1}{3}x^3 \ln(x^2) - \int \frac{2x^3}{3x} dx$
	$du = \frac{2x}{x^2} dx = \frac{2}{x} dx$	$v = \frac{1}{3}x^3$	$= \frac{1}{3}x^3 \ln(x^2) - \int \frac{2}{3}x^2 dx = \frac{1}{3}x^3 \ln(x^2) - \frac{2}{9}x^3 + c$

i) Parts then substitution:	$u = \arctan(2x)$	$dv = dx$	$\int \arctan(2x) dx = x \arctan(2x) - \int \frac{2x}{1+4x^2} dx$
	$du = \frac{2}{1+4x^2} dx$	$v = x$	
	$u = 4x^2$	$du = 4x dx$	So, $\int \frac{2x}{1+4x^2} dx = \frac{1}{4} \int \frac{1}{u} du = \frac{1}{4} \ln u  = \frac{1}{4} \ln(1+4x^2)$ , so
	$\frac{1}{8} du = 2x dx$		$\int \arctan(2x) dx = x \arctan(2x) - \frac{1}{4} \ln(1+4x^2) + c$

j)  $\int 2 \sec^3 x \tan x dx = 2 \int \sec^2 x (\sec x \tan x) dx = \frac{2}{3} \sec^3 x dx.$

k) Trig ID, then parts:  $\int x \sin^2 x dx = \int x [\frac{1}{2} - \frac{1}{2} \cos(2x)] dx = \int \frac{1}{2} x dx - \int \frac{1}{2} x \cos(2x) dx.$

$u = \frac{1}{2}x$	$dv = \cos(2x) dx$	$\int \frac{1}{2} x \cos(2x) dx = \frac{1}{4} x \sin(2x) - \int \frac{1}{4} \sin(2x) dx = \frac{1}{4} x \sin(2x) + \frac{1}{8} \cos(2x)$
$du = \frac{1}{2} dx$	$v = \frac{1}{2} \sin(2x)$	

Therefore:  $\int x \sin^2 x dx = \int \frac{1}{2} x dx - \int \frac{1}{2} x \cos(2x) dx = \frac{1}{4} x^2 - \frac{1}{4} x \sin(2x) - \frac{1}{8} \cos(2x).$


l) Reduction Formula:  $\int \tan^3 x dx = \frac{\tan^2 x}{2} - \int \tan x dx = \frac{\tan^2 x}{2} - \ln|\sec x| + c.$

6. First use parts:	$u = \arcsin x$	$dv = 2x dx$	$\int 2x \arcsin x dx = x^2 \arcsin x - \int \frac{x^2}{\sqrt{1-x^2}} dx$
	$du = \frac{1}{\sqrt{1-x^2}} dx$	$v = x^2$	Now use triangle substitution


$x = \sin \theta$	$dx = \cos \theta d\theta$	$\int \frac{x^2}{\sqrt{1-x^2}} dx = \int \frac{\sin^2 \theta \cos \theta}{\cos \theta} d\theta = \int \sin^2 \theta d\theta = \int \frac{1}{2} - \frac{1}{2} \cos(2\theta) d\theta$
$\sqrt{1-x^2} = \cos \theta$	$\theta = \arcsin x$	$= \frac{1}{2} \theta - \frac{1}{4} \sin(2\theta) = \frac{1}{2} \theta - \frac{1}{2} \sin(\theta) \cos(\theta) = \frac{1}{2} \arcsin x - \frac{1}{2} x \sqrt{1-x^2} + c$

So putting it all together  $\int 2x \arcsin x dx = x^2 \arcsin x - \frac{1}{2} \arcsin x + \frac{1}{2} x \sqrt{1-x^2} + c.$

7. a) In this case,  $u$  must correspond to the hypotenuse of the right triangle. (Why?) We have our choice of how to label the legs, one side  $a$  and the other  $\sqrt{u^2 - a^2}$ . With the selection below,  $u = a \sec \theta$ . What would  $u$  equal if we had let  $a$  be the side opposite  $\theta$ ?



- b) In this case,  $a$  must correspond to the hypotenuse of the right triangle. (Why?) We have a choice of how to label the legs, one side  $\sqrt{a^2 - u^2}$  and the other  $u$ . With the selection below,  $u = a \sin \theta$ , which is simpler than the other choice.



- c) Why must  $\sqrt{a^2 + u^2}$  must correspond to the hypotenuse of the right triangle? Why choose the  $u$  and  $a$  sides as follows rather than reverse their positions?



8. Where appropriate, use the triangles in the previous problem to help with the setup.

$$\text{a)} = \int \frac{\tan \theta}{\sec \theta} \sec \theta \tan \theta d\theta = \int \tan^2 \theta d\theta = \int (\sec^2 \theta - 1) d\theta = \tan \theta - \theta = \sqrt{x^2 - 1} - \arctan(\sqrt{x^2 - 1}) + c$$


$$\text{b)} = \int \frac{1}{6^3 \sec^3 \theta} 6 \sec^2 \theta d\theta = \frac{1}{36} \int \cos \theta d\theta = \frac{1}{36} \sin \theta = \frac{x}{36\sqrt{36 + x^2}} + c$$

$$\text{c)} = \int \frac{1}{4 \sec^2 \theta 2 \tan \theta} 2 \sec \theta \tan \theta d\theta = \int \frac{1}{4} \cos \theta d\theta = \frac{1}{4} \sin \theta = \frac{\sqrt{x^2 - 4}}{4x} + c$$

d) Substitution is easiest:  $u = 4 - x^2$  and  $du = -2x dx$ . So  $-\frac{1}{2} \int u^{1/2} du = -\frac{1}{3} u^{3/2} = -\frac{1}{3} (4 - x^2)^{3/2} + c$ .


e) Triangles:  $\int \frac{4}{4 - x^2} dx = \int \frac{4}{(\sqrt{4 - x^2})^2} dx = \int \frac{4}{(2 \cos \theta)^2} \cdot 2 \cos \theta d\theta = \int \frac{8 \cos \theta}{4 \cos^2 \theta} d\theta = 2 \int \sec \theta d\theta$   
 $= 2 \int \ln |\sec(\theta) + \tan \theta| d\theta = 2 \ln \left| \frac{2}{\sqrt{4 - x^2}} + \frac{x}{\sqrt{4 - x^2}} \right| + c = 2 \ln \left| \frac{2 + x}{\sqrt{4 - x^2}} \right| + c = \ln \left| \frac{(2 + x)^2}{4 - x^2} \right| + c = \ln \left| \frac{2 + x}{2 - x} \right| + c$

f)  $= \int \frac{8 \sin^3 \theta}{2 \cos \theta} 2 \cos \theta d\theta = \int 8 \sin^3 \theta d\theta = 8 \left[ -\frac{\sin^2 \theta \cos \theta}{3} + \frac{2}{3} \int \sin \theta d\theta \right] = 8 \left[ -\frac{\sin^2 \theta \cos \theta}{3} - \frac{2}{3} \cos \theta \right] + c = -\frac{x^2 \sqrt{4 - x^2}}{3} - \frac{8\sqrt{4 - x^2}}{3} + c = -\frac{1}{3} \sqrt{4 - x^2} (x^2 + 8) + c$

g)   $\frac{2x}{3} = \sin \theta \Rightarrow x = \frac{3}{2} \sin \theta$   
 $dx = \frac{3}{2} \cos \theta d\theta$   
 $\frac{\sqrt{4 - 9x^2}}{3} = \cos \theta \Rightarrow \sqrt{4 - 9x^2} = 3 \cos \theta$

$$\int \sqrt{9 - 4x^2} dx = \frac{9}{2} \int \cos^2 \theta d\theta = \frac{9}{2} \int \frac{1}{2} - \frac{1}{2} \cos 2\theta d\theta = \frac{9}{2} \left[ \frac{\theta}{2} - \frac{1}{4} \sin 2\theta \right] + c = \frac{9}{2} \left[ \frac{\theta}{2} - \frac{1}{2} \sin \theta \cos \theta \right] + c$$

$$= \frac{9}{2} \left[ \frac{\arcsin(2x/3)}{2} - \frac{1}{2} \cdot \frac{2x}{3} \cdot \frac{\sqrt{9 - 4x^2}}{3} \right] + c = \frac{9}{4} \arcsin(2x/3) - \frac{1}{2} x \sqrt{9 - 4x^2} + c$$

9.   $3x = \cos \theta \Rightarrow x = \frac{1}{3} \cos \theta$   
 $dx = -\frac{1}{3} \sin \theta d\theta$   
 $\sqrt{1 - 9x^2} = \sin \theta$  and  $9x^2 = \cos^2 \theta$

$$\int \frac{1}{9x^2 \sqrt{1 - 9x^2}} dx = \int \frac{-\frac{1}{3} \sin \theta}{\cos^2 \theta \sin \theta} d\theta = -\frac{1}{3} \int \sec^2 \theta d\theta = -\frac{1}{3} \tan \theta + c = -\frac{\sqrt{1 - 9x^2}}{9x} + c$$

10. Some Brief Answers to the More Integral Mix-Up Problem:

a) Substitution:  $u = 2x$ .  $\frac{1}{2} x e^{2x} - \frac{1}{4} e^{2x} + c$

b) Substitution:  $u = x^2 + 1$ .  $\frac{1}{2} e^{x^2 + 1} + c$

c) Parts twice (circular reasoning).  $\frac{1}{2} e^x \cos(x) + \frac{1}{2} e^x \sin(x) + c$

d) Parts twice.  $-x^2 \cos x + 2 \cos x + 2x \sin x + c$

e) Reduction:  $u = 2x$  so  $\frac{1}{2} du = dx$ .  $\int \sin^4(2x) dx = \frac{1}{2} \int \sin^4 u du$

$$= -\frac{1}{2} \left[ -\frac{\sin^3 u \cos u}{4} + \frac{3}{4} \int \sin^2 u du \right] = \frac{1}{2} \left[ -\frac{\sin^3 u \cos u}{4} + \frac{3}{4} \left( -\frac{1}{2} \sin u \cos u + \frac{1}{2} \int 1 du \right) \right]$$

$$= \frac{1}{2} \left[ -\frac{\sin^3 u \cos u}{4} + \frac{3}{4} \left( -\frac{1}{2} \sin u \cos u + \frac{u}{2} \right) \right] + c = -\frac{\sin^3 2x \cos 2x}{8} - \frac{3 \sin 2x \cos 2x}{16} + \frac{3x}{8} + c$$

f) Substitution:  $u = x^2$ .  $\frac{1}{2} \sin(x^2) + c$

g) Substitution:  $u = \ln x$ .  $\frac{(\ln x)^3}{3} + c$

h) First use  $\ln x^2 = 2 \ln x$ . Then use parts.  $2x \ln(x) - 2x + c$

i) First use  $\ln x^{1/2} = \frac{1}{2} \ln x$ . Then use parts.  $\frac{1}{2} x \ln(x) - \frac{1}{2} x + c$

j) Parts.  $\frac{1}{3} x^3 \left( \ln x - \frac{1}{3} \right) + c$

k) Half-angle.  $\frac{1}{2} x + \frac{1}{48} \sin(24x) + c$

- l) Ordinary:  $\ln |\sec x| + c$
- m) Parts then  $u$ -sub:  $\arctan x - \frac{1}{2} \ln(1 + x^2) + c$
- n) Parts twice.  $\frac{1}{3} \sin^3 x - \frac{1}{5} \sin^5 x + c$
- o) Substitution is easiest:  $u = 4 - x^2$  and  $du = -2x dx$ . So  $-\frac{1}{2} \int u^{1/2} du = -\frac{1}{3} u^{3/2} = -\frac{1}{3} (4 - x^2)^{3/2} + c$ .
- p) Let  $a = 3$ ,  $u = 2x$ ,  $du = 2dx$ , so  $\frac{1}{2} du = dx$ . Then

$$\int \frac{1}{\sqrt{9 - 4x^2}} dx = \frac{1}{2} \int \frac{1}{\sqrt{a^2 - u^2}} du = \frac{1}{2} \arcsin \frac{u}{a} + c = \frac{1}{2} \arcsin \frac{2x}{3} + c.$$