

## Functions

- a *functional relationship* between sets  $A$  and  $B$  associates exactly one element of  $B$  with each element of  $A$
- also called a *mapping* from set  $A$  to set  $B$
- a *function* expresses a functional relationship
  - written  $f: A \rightarrow B$ 
    - $f$  is a function from  $A$  to  $B$
    - $f$  maps  $A$  to  $B$
  - for  $a \in A$ ,  $f(a) \in B$  is the element of  $B$  that  $f$  associates with  $a$ 
    - $f(a)$  is the *value* of  $f$  at  $a$
  - note that while there is exactly one element of  $B$  associated with a given  $a$ , it is not required that it be unique i.e. it is not required that  $f(a_1) \neq f(a_2)$  for  $a_1 \neq a_2$
- $g \circ f$  is the *composition* of  $g$  and  $f$ :  $(g \circ f)(a) = g(f(a))$ 
  - to be valid, requires that  $f: A \rightarrow B$  and  $g: B \rightarrow C$

## Ordered $n$ -tuples

- $(a,b)$  is the *ordered pair* containing entities  $a$  and  $b$ 
  - if  $a \neq b$ ,  $(a,b)$  and  $(b,a)$  are different
  - $(a,b)$  and  $(c,d)$  are equal iff  $a = c$  and  $b = d$
- for sets  $A$  and  $B$ ,  $A \times B$  is the *cross product* (or *Cartesian product*)
 
$$A \times B = \{(a, b) \mid a \in A \wedge b \in B\}$$
  - contains every ordered pair containing an element of  $A$  and an element of  $B$
  - extends to *ordered triples*  $A \times B \times C$  and other *ordered  $n$ -tuples*
- the set  $\{(a,b) \in A \times B \mid a \in A \text{ and } b = f(a)\}$  is the *graph* of  $f$ 
  - a function can be specified by giving a set of ordered pairs
- $f((a,b))$  is more commonly written  $f(a,b)$

1. Let  $A = \{1, 2, 3, 4\}$  and let  $B = \{a, b, c\}$ . Find the sets  $A \times B$  and  $B \times A$ .

2. Let  $A$  be the set  $\{a, b, c, d\}$ . Let  $f$  be the function from  $A$  to  $A$  given by the set of ordered pairs  $\{(a, b), (b, b), (c, a), (d, c)\}$ , and let  $g$  be the function given by the set of ordered pairs  $\{(a, b), (b, c), (c, d), (d, d)\}$ . Find the set of ordered pairs for the composition  $g \circ f$ .

3. Let  $A = \{a, b, c\}$  and let  $B = \{0, 1\}$ . Find all possible functions from  $A$  to  $B$ . Give each function as a set of ordered pairs. (Hint: Every such function corresponds to one of the subsets of  $A$ .)

## More Terminology

for a function  $f: A \rightarrow B$

- $A$  is the *domain* of  $f$
- $B$  is the *range* of  $f$
- the *image* of  $f$  is  $\{f(a) \mid a \in A\}$ 
  - in some contexts, this is known as the *range*
- function  $f$  is *onto* (or *surjective*) if the image is equal to the range  $\forall b \in B (\exists a \in A (b = f(a)))$
- function  $f$  is *one-to-one* (or *injective*) if each element of the range is associated with at most one element of the domain  $\forall x \in A \forall y \in A (x \neq y \rightarrow f(x) \neq f(y))$   
 $\forall x \in A \forall y \in A (f(x) = f(y) \rightarrow x = y)$
- function  $f$  is *bijjective* if it is both one-to-one and onto

4. Consider the functions from  $\mathbb{Z}$  to  $\mathbb{Z}$  which are defined by the following formulas. Decide whether each function is onto and whether it is one-to-one; prove your answers.

a)  $f(n) = 2n$       b)  $g(n) = n + 1$       c)  $h(n) = n^2 + n + 1$

d)  $s(n) = \begin{cases} n/2, & \text{if } n \text{ is even} \\ (n+1)/2, & \text{if } n \text{ is odd} \end{cases}$